

Preprint: imsc-94/43

October 1994

On a nonstandard two-parametric quantum algebra and its connections with $U_{p,q}(gl(2))$ and $U_{p,q}(gl(1|1))$

R. Chakrabarti¹, R. Jagannathan² *

¹Department of Theoretical Physics, University of Madras, Guindy Campus, Madras-600025, India

²The Institute of Mathematical Sciences, C I T Campus, Tharamani, Madras-600113, India

Abstract. A quantum algebra $U_{p,q}(\zeta, H, X_{\pm})$ associated with a nonstandard R -matrix with two deformation parameters (p, q) is studied and, in particular, its universal $\mathcal{R}$ -matrix is derived using Reshetikhin's method. Explicit construction of the (p, q) -dependent nonstandard R -matrix is obtained through a coloured generalized boson realization of the universal $\mathcal{R}$ -matrix of the standard $U_{p,q}(gl(2))$ corresponding to a nongeneric case. General finite dimensional coloured representation of the universal $\mathcal{R}$ -matrix of $U_{p,q}(gl(2))$ is also derived. This representation, in nongeneric cases, becomes a source for various (p, q) -dependent nonstandard R -matrices. Superization of $U_{p,q}(\zeta, H, X_{\pm})$ leads to the super-Hopf algebra $U_{p,q}(gl(1|1))$. A contraction procedure then yields a (p, q) -deformed super-Heisenberg algebra $U_{p,q}(sh(1))$ and its universal $\mathcal{R}$ -matrix.

1. Introduction

The single parameter quantization of the universal enveloping algebra of a simple Lie algebra is well-known [1]. The Yang-Baxter equation (YBE), however, also admits nonstandard solutions [2-4] characterizing quasitriangular

*E-mail: jagan@imsc.ernet.in

Hopf algebras, which are not deformations of classical algebras. The nonstandard quantum algebra $U_q(\zeta, H, X_{\pm})$ associated with the Alexander-Conway solution of the YBE has been studied [2,3] and the relevant universal $\mathcal{R}$ -matrix has been obtained [4]. Using transmutation theory [5] it has been argued [4] that the superized $U_q(\zeta, H, X_{\pm})$ coincides with a super-Hopf algebra $U_q(gl(1|1))$. Moreover, using a general q -boson [6] realization of the Hopf algebra $U_q(gl(2))$, it has been observed [7] that the nonstandard R -matrix of $U_q(\zeta, H, X_{\pm})$ may be obtained from the universal $\mathcal{R}$ -matrix of $U_q(gl(2))$ in a nongeneric limit.

In another development, the constructions and representations of quantum algebras with multiple deformation parameters have been studied extensively [8-15]. For quasitriangular Hopf algebras, Reshetikhin [9] has developed a general formalism to introduce multiple deformation parameters. Following [9] the universal $\mathcal{R}$ -matrix of the quantum algebra $U_{p,q}(gl(2))$ with two independent parameters (p, q) has been obtained [15]. An identical procedure may be adopted to construct the universal $\mathcal{R}$ -matrix of the super-Hopf algebra $U_{p,q}(gl(1|1))$; this verifies the known result obtained by direct computation [13]. Here, we study a (p, q) -generalization, $U_{p,q}(\zeta, H, X_{\pm})$, of the Alexander-Conway algebra $U_q(\zeta, H, X_{\pm})$.

Following the prescription in [9] the universal $\mathcal{R}$ -matrix for the quasitriangular Hopf algebra $U_{p,q}(\zeta, H, X_{\pm})$ is obtained in section 2. The nonstandard Hopf algebra $U_{p,q}(\zeta, H, X_{\pm})$ has been previously considered in [16]. These authors, however, have not discussed the universal $\mathcal{R}$ -matrix for $U_{p,q}(\zeta, H, X_{\pm})$. Parallel to its single deformation parameter analogue, $U_{p,q}(\zeta, H, X_{\pm})$ exhibits close kinships with $U_{p,q}(gl(2))$ and $U_{p,q}(gl(1|1))$. In particular, the nonstandard R -matrix of $U_{p,q}(\zeta, H, X_{\pm})$ is obtained in section 3 using a coloured generalized boson representation of the universal $\mathcal{R}$ -matrix of $U_{p,q}(gl(2))$ in a nongeneric limit. A general recipe for realizing the finite dimensional nonstandard two-parametric coloured R -matrices associated with nongeneric representations of $U_{p,q}(gl(2))$ is also presented. In section 4, a map connect-

ing the two-parametric quantum algebras $U_{p,q}(\zeta, H, X_{\pm})$ and $U_{p,q}(gl(1|1))$ via the superization procedure is described. A contraction procedure is used in section 5 to extract a (p, q) -deformed quasitriangular super-Heisenberg algebra $U_{p,q}(sh(1))$. We conclude in section 6.

2. Quasitriangular Hopf algebra $U_{p,q}(\zeta, H, X_{\pm})$

We study the Hopf algebra associated with the two-parametric nonstandard solution [16] of the YBE

$$R = \begin{pmatrix} Q^{-1} & 0 & 0 & 0 \\ 0 & \lambda^{-1} & 0 & 0 \\ 0 & \sigma & \lambda & 0 \\ 0 & 0 & 0 & -Q \end{pmatrix}, \quad \sigma = Q^{-1} - Q. \quad (2.1)$$

The defining relation of the quantum inverse scattering method [17]

$$R(T \otimes \mathbb{1})(\mathbb{1} \otimes T) = (\mathbb{1} \otimes T)(T \otimes \mathbb{1})R, \quad (2.2)$$

with the R -matrix as given by (2.1), describes a transfer matrix

$$T = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad (2.3)$$

whose elements obey the braiding relations

$$\begin{aligned} ab &= p^{-1}ba, & ac &= q^{-1}ca, & db &= -p^{-1}bd, & dc &= -q^{-1}cd, \\ p^{-1}bc &= q^{-1}cb, & ad - da &= (p^{-1} - q)bc, & b^2 &= 0, & c^2 &= 0, \end{aligned} \quad (2.4)$$

where

$$p = \lambda Q, \quad q = \lambda^{-1}Q. \quad (2.5)$$

A conjugate R -matrix

$$\tilde{R} = \left(R^{(+)}\right)^{-1} = \begin{pmatrix} Q & 0 & 0 & 0 \\ 0 & \lambda^{-1} & -\sigma & 0 \\ 0 & 0 & \lambda & 0 \\ 0 & 0 & 0 & -Q^{-1} \end{pmatrix} \quad (2.6)$$

also fits (2.2) with the elements of T obeying (2.4). The matrices $R^{(\pm)}$ are defined by

$$R^{(+)} = PRP, \quad R^{(-)} = R^{-1}, \quad (2.7)$$

where P is the permutation matrix, given by

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \quad (2.8)$$

If a and d are invertable, the elements $\{a, b, c, d, a^{-1}, d^{-1}\}$ generate a Hopf algebra $A_{p,q}(R)$ (or $Fun_{p,q}(R)$) whose coalgebraic structure readily follows. The coproduct(Δ), counit(ϵ) and antipode(S) maps are, respectively, given by

$$\begin{aligned} \Delta(T) &= T \dot{\otimes} T, \\ \Delta(a^{-1}) &= a^{-1} \otimes a^{-1} - a^{-1}ba^{-1} \otimes a^{-1}ca^{-1}, \\ \Delta(d^{-1}) &= d^{-1} \otimes d^{-1} - d^{-1}cd^{-1} \otimes d^{-1}bd^{-1}, \end{aligned} \quad (2.9)$$

$$\epsilon(T) = \mathbb{1}, \quad (2.10)$$

$$S(T) = T^{-1}, \quad S(a^{-1}) = a - bd^{-1}c, \quad S(d^{-1}) = d - ca^{-1}b, \quad (2.11)$$

where

$$T^{-1} = \begin{pmatrix} a^{-1} + a^{-1}bd^{-1}ca^{-1} & -a^{-1}bd^{-1} \\ -d^{-1}ca^{-1} & d^{-1} + d^{-1}ca^{-1}bd^{-1} \end{pmatrix} \quad (2.12)$$

and $\dot{\otimes}$ denotes the tensor product coupled with usual matrix multiplication. In the Hopf algebra $A_{p,q}(R)$ an invertable ‘group-like’ element D exists:

$$D = ad^{-1} - bd^{-1}cd^{-1}, \quad D^{-1} = da^{-1} - ba^{-1}ca^{-1}. \quad (2.13)$$

Using (2.4), the commutation relations for D follow:

$$[D, a] = 0, \quad [D, d] = 0, \quad \{D, b\} = 0, \quad \{D, c\} = 0. \quad (2.14)$$

The induced coalgebra maps for D are obtained from the relations (2.9-2.12):

$$\Delta(D) = D \otimes D, \quad \epsilon(D) = 1, \quad S(D) = D^{-1}. \quad (2.15)$$

Using the FRT-approach [17], the commutation relations for the generators $(\zeta, H, X_{\pm})$ of the Hopf algebra $U_{p,q}(\zeta, H, X_{\pm})$, dual to the algebra $A_{p,q}(R)$, are obtained from the relations

$$R^{(+)} \left(L^{(\varepsilon_1)} \otimes \mathbb{1} \right) \left(\mathbb{1} \otimes L^{(\varepsilon_2)} \right) = \left(\mathbb{1} \otimes L^{(\varepsilon_2)} \right) \left(L^{(\varepsilon_1)} \otimes \mathbb{1} \right) R^{(+)}, \quad (2.16)$$

where $(\varepsilon_1, \varepsilon_2) = (+, +), (-, -), (+, -)$ and

$$L^{(+)} = \begin{pmatrix} p^{-H}q^{-\zeta} & \sigma p^{-H-\frac{1}{2}}X_- \\ 0 & gp^{-H}q^{\zeta} \end{pmatrix}, \quad L^{(-)} = \begin{pmatrix} q^H p^{\zeta} & 0 \\ -\sigma g q^{H-\frac{1}{2}}X_+ & g q^H p^{-\zeta} \end{pmatrix}, \quad (2.17)$$

with $g = (-1)^{\zeta-H}$. The corresponding commutation relations read

$$\begin{aligned} X_{\pm}^2 &= 0, & [H, X_{\pm}] &= \pm X_{\pm}, & \{X_+, X_-\} &= [2\zeta], \\ [\zeta, X] &= 0 & \forall X &= H, X_{\pm}, \end{aligned} \quad (2.18)$$

where

$$[X] = \frac{Q^X - Q^{-X}}{Q - Q^{-1}}. \quad (2.19)$$

The comultiplication maps for the generators are

$$\begin{aligned} \Delta(X_{\pm}) &= X_{\pm} \otimes g^{\mp 1} Q^{\zeta} \lambda^{\pm \zeta} + Q^{-\zeta} \lambda^{\mp \zeta} \otimes X_{\pm}, \\ \Delta(H) &= H \otimes \mathbb{1} + \mathbb{1} \otimes H, \quad \Delta(\zeta) = \zeta \otimes \mathbb{1} + \mathbb{1} \otimes \zeta, \end{aligned} \quad (2.20)$$

which follow from the relations

$$\Delta(L^{(\pm)}) = L^{(\pm)} \dot{\otimes} L^{(\pm)}. \quad (2.21)$$

The counit and the antipode maps are given by

$$\epsilon(X) = 0, \quad \forall X = \zeta, H, X_{\pm}, \quad (2.22)$$

$$S(\zeta) = -\zeta, \quad S(H) = -H, \quad S(X_{\pm}) = g^{\pm 1} X_{\pm}. \quad (2.23)$$

In spite of the appearance of anticommutator in (2.18), the Hopf algebra $U_{p,q}(\zeta, H, X_{\pm})$ is bosonic as it follows the direct product rule

$$(A \otimes B)(C \otimes D) = AC \otimes BD, \quad \forall A, B, C, D \in U_{p,q}(\zeta, H, X_{\pm}). \quad (2.24)$$

For the finite dimensional faithful representation [3] of the algebra (2.18)

$$\begin{aligned} X_+ &= \begin{pmatrix} 0 & [z] \\ 0 & 0 \end{pmatrix}, \quad X_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad H = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \\ \zeta &= \frac{z}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \end{aligned} \quad (2.25)$$

with $z \in \mathbb{C}$, the duality relations between the Hopf algebras $A_{p,q}(R)$ and $U_{p,q}(\zeta, H, X_{\pm})$ assume the form

$$\langle L^{(\pm)}, T \rangle = R^{(\pm)}, \quad (2.26)$$

for the choice $z = 1$.

We now discuss the quasitriangular character of $U_{p,q}(\zeta, H, X_{\pm})$ containing a group-like element g , that is, now and henceforth, assumed to satisfy $g^2 = 1$. For a quasitriangular Hopf algebra $\mathcal{U}$, the universal $\mathcal{R}$ -matrix ($\in \mathcal{U} \otimes \mathcal{U}$) satisfies the relations

$$\begin{aligned} \tau \circ \Delta(X) &= \mathcal{R} \Delta(X) \mathcal{R}^{-1}, & \tau \circ (X \otimes Y) &= Y \otimes X, \quad \forall X, Y \in \mathcal{U}, \\ (\Delta \otimes \text{id}) \mathcal{R} &= \mathcal{R}_{13} \mathcal{R}_{23}, & (\text{id} \otimes \Delta) \mathcal{R} &= \mathcal{R}_{13} \mathcal{R}_{12}, \\ (\epsilon \otimes \text{id}) &= (\text{id} \otimes \epsilon) \mathcal{R} = \mathbb{1}, \\ (S \otimes \text{id}) \mathcal{R} &= \mathcal{R}^{-1}, & (\text{id} \otimes S) \mathcal{R}^{-1} &= \mathcal{R}, \end{aligned} \quad (2.27)$$

where the subscripts in $\mathcal{R}_{ij}$ indicate the embedding of $\mathcal{R}$ in $\mathcal{U}^{\otimes 3}$. The explicit expression for the universal $\mathcal{R}$ -matrix of $U_Q(\zeta, H, X_{\pm})$ is [4]

$$\mathcal{R}_{\lambda=1} = (-1)^{(\zeta-H) \otimes (\zeta-H)} Q^{2(\zeta \otimes H + H \otimes \zeta)} \left(\mathbb{1} \otimes \mathbb{1} + \sigma Q^{\zeta} X_+ \otimes g Q^{-\zeta} X_- \right). \quad (2.28)$$

Employing the Reshetikhin procedure [9] we obtain the universal $\mathcal{R}$ -matrix for $U_{p,q}(\zeta, H, X_{\pm})$. To this end, we note that the coproduct relations (2.20) for the generators of $U_{p,q}(\zeta, H, X_{\pm})$ and the corresponding relations for the generators of $U_Q(\zeta, H, X_{\pm})$, obtained in the limit $\lambda = 1$, are related by a similarity transformation

$$\Delta(X) = F(\Delta_{\lambda=1}(X)) F^{-1}, \quad \forall X = \zeta, H, X_{\pm}, \quad (2.29)$$

where

$$F = \lambda^{(H \otimes \zeta - \zeta \otimes H)}. \quad (2.30)$$

Then, the procedure in [9] yields the universal $\mathcal{R}$ -matrix for $U_{p,q}(\zeta, H, X_{\pm})$

$$\mathcal{R} = F^{-1} \mathcal{R}_{\lambda=1} F^{-1} \quad (2.31)$$

which, by construction, satisfies the required relations (2.27). Explicitly, the universal $\mathcal{R}$ -matrix of $U_{p,q}(\zeta, H, X_{\pm})$ reads

$$\begin{aligned} \mathcal{R} = & (-1)^{(\zeta-H) \otimes (\zeta-H)} Q^{2(\zeta \otimes H + H \otimes \zeta)} \lambda^{2(\zeta \otimes H - H \otimes \zeta)} \\ & \times \left(\mathbb{1} \otimes \mathbb{1} + \sigma Q^{\zeta} \lambda^{\zeta} X_+ \otimes g Q^{-\zeta} \lambda^{\zeta} X_- \right). \end{aligned} \quad (2.32)$$

For the representation (2.25) with the choice $z = 1$, $\mathcal{R}$ in (2.32) reduces to the matrix $\tilde{R}$ in (2.6), thus expressing an aspect of the duality of $A_{p,q}(R)$ and $U_{p,q}(\zeta, H, X_{\pm})$.

3. Coloured R -matrices associated with nongeneric representations of $U_{p,q}(gl(2))$

After completing the above construction of the dual Hopf algebras corresponding to the nonstandard solution (2.1) of the YBE, we now relate the R -matrix (2.1) to the representations of $U_{p,q}(gl(2))$ for nongeneric values of Q , namely, roots of unity. Here, we closely follow the treatment of $U_q(gl(2))$ by Ge et al. [7]. These authors have constructed a general parameter dependent finite dimensional q -boson realization of the universal $\mathcal{R}$ -matrix of $U_q(gl(2))$; thereby generating the nonstandard R -matrices at q , a root of unity. When different parameters are chosen for different components of the representation module, the resulting finite dimensional R -matrices are said to be coloured and they obey a coloured YBE [7].

In the standard Hopf algebraic structure of $U_{p,q}(gl(2))$ (see, e.g., [15]) the commutation relations between the generators are

$$[J_0, J_\pm] = \pm J_\pm, \quad [J_+, J_-] = [2J_0], \quad [Z, X] = 0, \quad \forall X \in (J_0, J_\pm), \quad (3.1)$$

and the comultiplication rules are

$$\begin{aligned} \Delta(X_\pm) &= J_\pm \otimes Q^{J_0} \lambda^{\pm Z} + Q^{-J_0} \lambda^{\mp Z} \otimes J_\pm, \\ \Delta(J_0) &= J_0 \otimes \mathbb{1} + \mathbb{1} \otimes J_0, \quad \Delta(Z) = Z \otimes \mathbb{1} + \mathbb{1} \otimes Z. \end{aligned} \quad (3.2)$$

The universal $\mathcal{R}$ -matrix of $U_{p,q}(gl(2))$ reads [15]

$$\begin{aligned} \bar{\mathcal{R}} &= Q^{2(J_0 \otimes J_0)} \lambda^{2(Z \otimes J_0 - J_0 \otimes Z)} \\ &\times \sum_{n=0}^{\infty} \frac{(1 - Q^{-2})^n}{[n]!} Q^{\frac{1}{2}n(n-1)} \left(Q^{J_0} \lambda^Z J_+ \otimes Q^{-J_0} \lambda^Z J_- \right)^n, \end{aligned} \quad (3.3)$$

where $[n]! = [n][n-1] \dots [2][1]$.

The generators of the deformed boson algebra [6] satisfy

$$[N, a_\pm] = \pm a_\pm, \quad a_+ a_- = [N], \quad a_- a_+ = [N+1], \quad (3.4)$$

and the map [7]

$$J_+ = a_+[\omega - N], \quad J_- = a_-, \quad J_0 = N - \frac{\omega}{2}, \quad \omega \in \mathbb{C}, \quad (3.5)$$

provides an infinite dimensional representation of the algebra (3.1) in the Fock space $\mathcal{F}_z \left\{ |mz\rangle = a_+^m |0z\rangle \mid a_- |0z\rangle = 0, N|0z\rangle = 0, m \in \mathbb{Z}^+, z \in \mathbb{C} \right\}$:

$$\begin{aligned} J_+ |mz\rangle &= [\omega - m] |m+1z\rangle, \quad J_- |mz\rangle = [m] |m-1z\rangle, \\ J_0 |mz\rangle &= \left(m - \frac{\omega}{2} \right) |mz\rangle, \quad Z |mz\rangle = z |mz\rangle. \end{aligned} \quad (3.6)$$

The parameter ω is called colour and provides the key to obtain the non-standard R -matrix (2.1) starting from the universal $\mathcal{R}$ -matrix of $U_{p,q}(gl(2))$, namely $\bar{\mathcal{R}}$ in (3.3).

For the nongeneric cases $\{Q^n = \pm 1, n \in \mathbb{Z}^+\}$ the identity $[\alpha n] = 0$ holds for $\alpha \in \mathbb{Z}^+$ suggesting the existence of an extremal vector $\{|\alpha n z\rangle \mid J_-|\alpha n z\rangle = 0\}$. The corresponding invariant subspace that renders the representation (3.6) reducible is $\mathcal{V}_{\alpha z}\{|\alpha n + m z\rangle \mid m \in \mathbb{Z}^+\}$. Then, on the quotient space $V_{Jz} = \mathcal{F}_z/\mathcal{V}_{\alpha z}\{|JMz\rangle (= |mz\rangle) \mid m = 0, 1, \dots, (\alpha n - 1) = 2J, M = m - J\}$ a finite dimensional representation for algebra (3.1) holds:

$$\begin{aligned} J_+|JMz\rangle &= [\omega - J - M]|J(M+1)z\rangle\theta(J-1-M), \\ J_-|JMz\rangle &= [J+M]|J(M-1)z\rangle, \\ J_0|JMz\rangle &= \left(J+M-\frac{\omega}{2}\right)|JMz\rangle, \\ Z|JMz\rangle &= z|JMz\rangle, \end{aligned} \tag{3.7}$$

where $\theta(x) = 1$ (0) for $x \geq 0$ (< 0). For $\alpha = 1$, the representation (3.7) is irreducible and for $\alpha \geq 2$, it is indecomposable [7]. Now, using the representation (3.7), $\bar{\mathcal{R}}$ in (3.3) may be written in the matrix form

$$\bar{\mathcal{R}}|J_1 M_1 z_1\rangle \otimes |J_2 M_2 z_2\rangle = \sum_{M'_1 M'_2} \left(\bar{\mathcal{R}}^{J_1 z_1 \omega_1, J_2 z_2 \omega_2}\right)_{M_1 M_2}^{M'_1 M'_2} |J_1 M'_1 z_1\rangle \otimes |J_2 M'_2 z_2\rangle \tag{3.8}$$

where different representations and colour parameters are chosen in the two sectors of the tensor product space. Explicitly we have

$$\begin{aligned} &\left(\bar{\mathcal{R}}^{J_1 z_1 \omega_1, J_2 z_2 \omega_2}\right)_{M_1 M_2}^{M'_1 M'_2} \\ &= Q^{2\left(J_1+M'_1-\frac{\omega_1}{2}\right)\left(J_2+M'_2-\frac{\omega_2}{2}\right)} \lambda^{2\left(z_1\left(J_2+M'_2-\frac{\omega_2}{2}\right)-z_2\left(J_1+M'_1-\frac{\omega_1}{2}\right)\right)} \\ &\quad \left\{ \sum_{n=0}^{\infty} \frac{(1-Q^{-2})^n}{[n]!} Q^{-\frac{1}{2}n(n-1)} Q^{n\left(J_1-J_2+M'_1-M'_2-\frac{1}{2}(\omega_1-\omega_2)\right)} \lambda^{n(z_1+z_2)} \right\} \end{aligned}$$

$$\Pi_{l=1}^n [\omega_1 - J_1 - M'_1 + l] [J_2 + M'_2 + l] \delta_{M'_1+n}^{M'_1} \delta_{M'_2-n}^{M'_2} \Big\} . \quad (3.9)$$

It should be noted that the presence of the second deformation parameter λ in (3.9) will produce many new R -matrices. For nongeneric values of Q , the matrix representation (3.9) acts as a source for obtaining the finite dimensional two parametric(p, q) nonstandard coloured R -matrices. Let us consider the simplest example ($J_1 = J_2 = \frac{1}{2}$, $z_1 = z_2 = z$) with different colour parameters in the two sectors of the Hilbert space. For $Q^2 = -1$, we get

$$\bar{\mathcal{R}}^{\frac{1}{2}z\omega_1, \frac{1}{2}z\omega_2} = c \begin{pmatrix} t_1 & 0 & 0 & 0 \\ 0 & s^{-1} & w & 0 \\ 0 & 0 & st_1 t_2^{-1} & 0 \\ 0 & 0 & 0 & -t_2^{-1} \end{pmatrix}, \quad (3.10)$$

where $c = Q^{-\omega_2(1-\frac{\omega_1}{2})} \lambda^{z(\omega_1-\omega_2)}$, $t_1 = -Q^{-\omega_1}$, $t_2 = -Q^{-\omega_2}$, $s = \lambda^{2z}$ and $w = (t_1 - t_1^{-1}) t_1^{\frac{1}{2}} t_2^{-\frac{1}{2}}$. This is an example of coloured R -matrix and may be viewed as a generalization of the result obtained in [7]. When $\omega_1 = \omega_2 = \omega$ the R -matrix (3.10) reduces to

$$\bar{\mathcal{R}}^{\frac{1}{2}z\omega, \frac{1}{2}z\omega} \sim \begin{pmatrix} t & 0 & 0 & 0 \\ 0 & s^{-1} & t - t^{-1} & 0 \\ 0 & 0 & s & 0 \\ 0 & 0 & 0 & -t^{-1} \end{pmatrix}. \quad (3.11)$$

Apart from a scale factor, the matrix in (3.11) agrees with the nonstandard R -matrix $\tilde{R}$ in (2.6) after the replacement $Q \rightarrow t$, $\lambda \rightarrow s$. This completes our discussion of the connection between the nonstandard two-parameter R -matrix (2.1) with the universal $\mathcal{R}$ -matrix of $U_{p,q}(gl(2))$.

4. Superization of $U_{p,q}(\zeta, H, X_{\pm})$

Using a superization procedure [4] we now discuss the connection between $U_{p,q}(\zeta, H, X_{\pm})$ and the super-Hopf algebra $U_{p,q}(gl(1|1))$. In [4] it is argued that if $\mathcal{H}$ is a Hopf algebra containing a group-like element g such that g^2

$= 1$, then, there exists a super-Hopf algebra $\hat{\mathcal{H}}$ with identical algebraic and counit structures while the coproduct, antipode and the universal R -matrix of $\mathcal{H}$, $\{\Delta(h) (= \sum_k x_k \otimes y_k), S(h), \mathcal{R} (= \sum_k X_k \otimes Y_k) \mid (h, x_k, y_k, X_k, Y_k) \in \mathcal{H}\}$, are modified to the corresponding quantities of $\hat{\mathcal{H}}$:

$$\begin{aligned}\Delta(\hat{h}) &= \sum_k x_k \otimes g^{\deg(x_k)} y_k, \quad S(\hat{h}) = S(h) g^{\deg(h)}, \\ \hat{\mathcal{R}} &= \mathcal{R}_g \sum_k X_k \otimes g^{\deg(X_k)} Y_k, \\ \mathcal{R}_g &= \frac{1}{2}(\mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes g + g \otimes \mathbb{1} - g \otimes g). \quad (4.1)\end{aligned}$$

The map $\{h \rightarrow \hat{h} \mid h \in \mathcal{H}, \hat{h} \in \hat{\mathcal{H}}\}$ preserves the algebraic structure and it is understood that, in the right hand side of (4.1) the elements of $\mathcal{H}$, after simplification, are mapped to their hatted superpartners in $\hat{\mathcal{H}}$. The $\deg(h)$, $\forall h \in \mathcal{H}$, is given by $ghg^{-1} = \deg(h)h$. We exhibit these properties in (p, q) -deformed examples.

The R -matrix

$$\hat{R} = \begin{pmatrix} Q^{-1} & 0 & 0 & 0 \\ 0 & \lambda^{-1} & 0 & 0 \\ 0 & \sigma & \lambda & 0 \\ 0 & 0 & 0 & Q \end{pmatrix} \quad (4.2)$$

satisfies the super-YBE and is known [11] to describe the ‘function’ Hopf algebra $Fun_{p,q}(GL(1|1))$ (or $GL_{p,q}(1|1)$). The universal $\mathcal{R}$ -matrix $\hat{\mathcal{R}}$ for the dually paired enveloping algebra $U_{p,q}(gl(1|1))$ [12] has been obtained by direct computation [13]. We can derive this $\hat{\mathcal{R}}$ by applying Reshetikhin’s technique [9] for introducing multiple deformation parameters as follows. The commutation relations for the generators $(\hat{\zeta}, \hat{H}, \hat{X}_{\pm})$ of $U_{p,q}(gl(1|1))$ read

$$\begin{aligned}\hat{X}_{\pm}^2 &= 0, \quad [\hat{H}, \hat{X}_{\pm}] = \pm \hat{X}_{\pm}, \quad \{\hat{X}_+, \hat{X}_-\} = [2\hat{\zeta}], \\ [\hat{\zeta}, X] &= 0, \quad \forall X \in (\hat{H}, \hat{X}_{\pm}), \quad (4.3)\end{aligned}$$

and the coalgebraic structure is

$$\begin{aligned}
\Delta(\hat{X}_\pm) &= \hat{X}_\pm \otimes Q^{\hat{\zeta}} \lambda^{\pm \hat{\zeta}} + Q^{-\hat{\zeta}} \lambda^{\mp \hat{\zeta}} \otimes \hat{X}_\pm, \\
\Delta(\hat{H}) &= \hat{H} \otimes \mathbb{1} + \mathbb{1} \otimes \hat{H}, \quad \Delta(\hat{\zeta}) = \hat{\zeta} \otimes \mathbb{1} + \mathbb{1} \otimes \hat{\zeta}, \\
\epsilon(X) &= 0, \quad S(X) = -X, \quad \forall X \in (\hat{\zeta}, \hat{H}, \hat{X}_\pm). \quad (4.4)
\end{aligned}$$

The coproduct relations (4.4) reveal that a structure similar to (2.29) holds with the choice

$$\hat{F} = \lambda^{(\hat{H} \otimes \hat{\zeta} - \hat{\zeta} \otimes \hat{H})}. \quad (4.5)$$

Now, using the known universal $\mathcal{R}$ -matrix of $U_Q(gl(1|1))$ [4]

$$\hat{\mathcal{R}}_{\lambda=1} = Q^{2(\hat{\zeta} \otimes \hat{H} + \hat{H} \otimes \hat{\zeta})} (\mathbb{1} \otimes \mathbb{1} + \sigma Q^{\hat{\zeta}} \hat{X}_+ \otimes Q^{-\hat{\zeta}} \hat{X}_-) \quad (4.6)$$

and the prescription (2.31), we obtain the universal $\mathcal{R}$ -matrix for $U_{p,q}(gl(1|1))$ as

$$\hat{\mathcal{R}} = Q^{2(\hat{\zeta} \otimes \hat{H} + \hat{H} \otimes \hat{\zeta})} \lambda^{2(\hat{\zeta} \otimes \hat{H} - \hat{H} \otimes \hat{\zeta})} (\mathbb{1} \otimes \mathbb{1} + \sigma Q^{\hat{\zeta}} \lambda^{\hat{\zeta}} \hat{X}_+ \otimes Q^{-\hat{\zeta}} \lambda^{\hat{\zeta}} \hat{X}_-) \quad (4.7)$$

which satisfies the super-YBE with the composition rule for the graded operators $(A, B, C, D) \in U_{p,q}(gl(1|1))$

$$(A \otimes B)(C \otimes D) = (-1)^{\deg(B)\deg(C)}(AC \otimes BD). \quad (4.8)$$

Our expression for $\hat{\mathcal{R}}$ in (4.7) is seen to agree with the result obtained in [13] by direct computation.

Comparing (2.18) and (2.22) with (4.3) and (4.4), while introducing the map $(\zeta, H, X_\pm) \rightarrow (\hat{\zeta}, \hat{H}, \hat{X}_\pm)$, it follows that the commutation relations and the counit maps for the super-Hopf algebra $U_{p,q}(gl(1|1))$ are identical to those of $U_{p,q}(\zeta, H, X_\pm)$. The coproduct (4.4), antipode (4.4) and the universal

$\mathcal{R}$ -matrix (4.7) for $U_{p,q}(gl(1|1))$ follow from the analogous formulae, (2.20), (2.23) and (2.32), for $U_{p,q}(\hat{\zeta}, \hat{H}, \hat{X}_{\pm})$ according to the prescription (4.1). The quantity $(-1)^{(\zeta-H)\otimes(\zeta-H)}\mathcal{R}_g$ is central in nature as may be observed by direct computation. It may, therefore, be dropped while mapping $\mathcal{R}$ to $\hat{\mathcal{R}}$.

5. A (p, q) -deformed super-Heisenberg algebra

Finally, we use the contraction technique à la Celeghini et al. [18,19] to extract a two-parametric deformed super-Heisenberg algebra $U_{p,q}(sh(1))$ as a limiting case of $U_{p,q}(gl(1|1))$. To this end, let us scale the generators and the quantization parameters as

$$\hat{H} = \frac{1}{2\varepsilon}\mathbf{h} - N, \quad \hat{\zeta} = \frac{\xi}{2\varepsilon}, \quad \hat{X}_{\mp} = \frac{1}{\sqrt{\varepsilon}}c_{\pm}, \quad Q = e^{\varepsilon\Omega}, \quad \lambda = e^{\varepsilon\nu}. \quad (5.1)$$

The commutation relations and the coproduct rules for $U_{p,q}(sh(1))$, with $\{\xi, \mathbf{h}, N, c_{\pm}\}$ as the generators, are obtained by studying the $\varepsilon \rightarrow 0$ limit of the corresponding structures of the algebra $U_{p,q}(gl(1|1))$ in (4.3) and (4.4) respectively. The result is

$$\begin{aligned} c_{\pm}^2 &= 0, & [N, c_{\pm}] &= \pm c_{\pm}, & \{c_+, c_-\} &= \Omega^{-1} \sinh \Omega \xi, \\ [\mathbf{h}, X] &= 0, & [\xi, X] &= 0, & \forall X \in U_{p,q}(sh(1)), \end{aligned} \quad (5.2)$$

and

$$\begin{aligned} \Delta(c_{\pm}) &= c_{\pm} \otimes e^{\frac{1}{2}(\Omega \mp \nu)\xi} + e^{-\frac{1}{2}(\Omega \mp \nu)\xi} \otimes c_{\pm}, \\ \Delta(\xi) &= \xi \otimes \mathbb{1} + \mathbb{1} \otimes \xi, \quad \Delta(\mathbf{h}) = \mathbf{h} \otimes \mathbb{1} + \mathbb{1} \otimes \mathbf{h}, \\ \Delta(N) &= N \otimes \mathbb{1} + \mathbb{1} \otimes N. \end{aligned} \quad (5.3)$$

In the contraction limit, the universal $\mathcal{R}$ -matrix (4.7) of $U_{p,q}(gl(1|1))$ yields, after the removal of a constant singular factor, the universal $\mathcal{R}$ -matrix of $U_{p,q}(sh(1))$:

$$\mathcal{R}^{sh} = e^{\nu(N \otimes \xi - \xi \otimes N) - \Omega(N \otimes \xi + \xi \otimes N)} \left(\mathbb{1} \otimes \mathbb{1} - 2\Omega e^{\frac{1}{2}(\Omega + \nu)\xi} c_- \otimes e^{-\frac{1}{2}(\Omega - \nu)\xi} c_+ \right). \quad (5.4)$$

A spectral parameter dependent $\mathcal{R}$ -matrix may be obtained from the universal $\mathcal{R}$ -matrix (5.4). To this end, following [18], we define the map

$$T_x c_{\pm} = x^{\mp 1} c_{\pm}, \quad T_x \mathbf{h} = \mathbf{h}, \quad T_x \xi = \xi, \quad T_x N = N, \quad (5.5)$$

and let

$$\begin{aligned} \mathcal{R}^{sh}(x) &= (T_x \otimes \mathbb{1}) \mathcal{R}^{sh} \\ &= e^{\nu(N \otimes \xi - \xi \otimes N) - \Omega(N \otimes \xi + \xi \otimes N)} \\ &\quad \times \left(\mathbb{1} \otimes \mathbb{1} - 2\Omega x e^{\frac{1}{2}(\Omega + \nu)\xi} c_- \otimes e^{-\frac{1}{2}(\Omega - \nu)\xi} c_+ \right). \end{aligned} \quad (5.6)$$

A direct calculation then proves that the matrix $\mathcal{R}^{sh}(x)$ satisfies the spectral parameter dependent YBE

$$\mathcal{R}_{12}^{sh}(x) \mathcal{R}_{13}^{sh}(xy) \mathcal{R}_{23}^{sh}(y) = \mathcal{R}_{23}^{sh}(y) \mathcal{R}_{13}^{sh}(xy) \mathcal{R}_{12}^{sh}(x). \quad (5.7)$$

6. Conclusion

To conclude, we have studied the dually paired Hopf algebras $A_{p,q}(R)$ and $U_{p,q}(\zeta, H, X_{\pm})$ associated with the nonstandard R -matrix (2.1) involving two independent parameters p and q . The universal $\mathcal{R}$ -matrix of $U_{p,q}(\zeta, H, X_{\pm})$ has been obtained. We have demonstrated an explicit construction of the nonstandard R -matrix through a coloured generalized boson representation of the universal $\mathcal{R}$ -matrix of $U_{p,q}(gl(2))$ corresponding to the nongeneric case $pq = Q^2 = -1$. In this example, by choosing different colour parameters for the two sectors of nongeneric representations of $U_{p,q}(gl(2))$ with a dimension 2, we have obtained a two-parametric coloured R -matrix (3.10), which

satisfies a coloured YBE. More importantly, the finite dimensional representation (3.9) of the $\mathcal{R}$ -matrix gives a recipe for obtaining nonstandard two-parametric coloured R -matrices for nongeneric values of Q . Superization process describes a map between $U_{p,q}(\zeta, H, X_{\pm})$ and $U_{p,q}(gl(1|1))$. A contraction procedure has been used to obtain a (p, q) -deformed quasitriangular super-Heisenberg algebra $U_{p,q}(sh(1))$.

References

1. V.G. Drinfeld: Proc. ICM-86 (Berkeley) 1987; M. Jimbo: Lett. Math. Phys. 10 (1985) 63
2. H.C. Lee, M. Couture, N.C. Schmeing: ‘Connected link polynomials’, Chalk-river preprint 1988; M.L. Ge, L.Y. Wang, K. Xue, Y.S. Wu: Int. J. Mod. Phys. A 4 (1989) 3351
3. N. Jing, M.L. Ge, Y.S. Wu: Lett. Math. Phys. 21 (1991) 193
4. S. Majid, M.J. Rodriguez-Plaza: ‘Universal $\mathcal{R}$ -matrix for a nonstandard quantum group and superization’, Preprint DAMTP 91/47
5. S. Majid: ‘Transmutation theory and rank for quantum braided groups’ Preprint DAMTP 91/10
6. A.J. Macfarlane: J. Phys. A: Math. Gen. 22 (1989) 4551; L.C. Biedenharn: J. Phys. A: Math. Gen. 22 (1989) L873; C.P. Sun, H.C. Fu: J. Phys. A: Math. Gen. 22 (1989) L983; T. Hayashi: Commun. Math. Phys. 127 (1990) 129
7. M.L. Ge, C.P. Sun, K. Xue: Int. J. Mod. Phys. A 7 (1992) 6609
8. P.P. Kulish: Zap. Nauch. Semin. LOMI 180 (1990) 89; E.E. Demidov, Yu.I. Manin, E.E. Mukhin, D.V. Zhadanovich: Prog. Theor. Phys. Suppl. 102 (1990) 203; A. Sudbery: J. Phys. A: Math. Gen. 23 (1990) L697; M. Takeuchi: Proc. Japan Acad. 66 (1990) 112
9. N.Yu. Reshetikhin: Lett. Math. Phys. 20 (1990) 331
10. A. Schirmacher, J. Wess, B. Zumino: Z. Phys. C: Particles and Fields 49 (1991) 317; A. Schirmacher: J. Phys. A: Math. Gen. 24 (1991) L1249; C. Burdick, L. Hlavaty: J. Phys. A: Math. Gen. 24 (1991) L165

11. R. Chakrabarti, R. Jagannathan: J. Phys. A: Math. Gen. 24 (1991) 5683
12. L. Dobrowski, L.Y. Wang: Phys. Lett. B226 (1991) 51
13. C. Burdick, P. Hellinger: J. Phys. A: Math. Gen. 25 (1992) L1023
14. V.K. Dobrev: J. Geom. Phys. 11 (1993) 367; C. Fronsdal, A. Galindo: Lett. Math. Phys. 27 (1993) 59; R. Barbier, J. Meyer, M. Kibler: J. Phys. G: Nucl. Part. 20 (1994) L13
15. R. Chakrabarti, R. Jagannathan: J. Phys. A: Math. Gen. 27 (1994) 2023
16. M. Bednar, C. Burdick, M. Couture, L. Hlavaty: J. Phys. A: Math. Gen. 25 (1992) L341
17. N.Yu. Reshetikhin, L.A. Takhtajan, L.D. Faddeev: Leningrad Math. J. 1 (1990) 193
18. E. Celeghini, R. Giachetti, E. Sorace, M. Tarlini: J. Math. Phys. 31 (1991) 2548
19. E. Celeghini, R. Giachetti, E. Sorace, M. Tarlini: J. Math. Phys. 32 (1992) 1115

Note added: We are thankful to Prof. V.K. Dobrev for bringing our attention to the papers: (1) V.K. Dobrev, "Introduction to Quantum Groups", Göttingen University preprint, (April 1991), to appear in: *Proc. 22nd Ann. Iranian Math. Conf.* (March 1991, Mashhad) and (2) V.K. Dobrev, *J. Math. Phys.* 33 (1992) 3419, where the structure of $U_{p,q}(gl(2))$ dual to $GL_{p,q}(2)$ was established earlier by him (see Refs. in [14]) using an approach independent of Schirrmacher, Wess and Zumino [10]. It may also be noted that the work of Fronsdal and Galindo (see Refs. in [14]) on the $U_{p,q}(gl(2)) \longrightarrow GL_{p,q}(2)$ exponential relationship is based on a different approach.